


Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1




1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526

J. L. Chen, L. Dai, P. Zhang, W. Chen, G. S. Cai, X. W. Qi, M. Z. Hu, B. Du, Q. F. Pang. Methylene blue attenuates acute liver injury induced by paraquat in rats. <i>Int Immunopharmacol.</i> 2015. 28:808-12
M. Ren, Y. M. Wang, J. Zhao, J. Zhao, Z. M. Zhao, T. F. Zhang, J. He, S. P. Ren, S. Q. Peng. Metallothioneins attenuate paraquat-induced acute lung injury in mice through the mechanisms of anti-oxidation and anti-apoptosis. <i>Food Chem Toxicol.</i> 2014. 73:140-7
B. J. Day, J. D. Crapo. A metalloporphyrin superoxide dismutase mimetic protects against paraquat-induced lung injury in vivo. <i>Toxicol Appl Pharmacol.</i> 1996. 140:94-100
J. S. Bus, S. Z. Cagen, M. Olgaard, J. E. Gibson. A mechanism of paraquat toxicity in mice and rats. <i>Toxicol Appl Pharmacol.</i> 1976. 35:501-13
T. Fukushima, K. Yamada, N. Hojo, A. Isobe, K. Shiwaku, Y. Yamane. Mechanism of cytotoxicity of paraquat. III. The effects of acute paraquat exposure on the electron transport system in rat mitochondria. <i>Exp Toxicol Pathol.</i> 1994. 46:437-41
K. Yamada, T. Fukushima. Mechanism of cytotoxicity of paraquat. II. Organ specificity of paraquat-stimulated lipid peroxidation in the inner membrane of mitochondria. <i>Exp Toxicol Pathol.</i> 1993. 45:375-80
I. Ahmad, S. Shukla, A. Kumar, B. K. Singh, D. K. Patel, H. P. Pandey, C. Singh. Maneb and paraquat-induced modulation of toxicant responsive genes in the rat liver: comparison with polymorphonuclear leukocytes. <i>Chem Biol Interact.</i> 2010. 188:566-79
W. D. Huang, J. Z. Wang, Y. Q. Lu, Y. M. Di, J. K. Jiang, Q. Zhang. Lysine acetylsalicylate ameliorates lung injury in rats acutely exposed to paraquat. <i>Chin Med J (Engl).</i> 2011. 124:2496-501
J. W. Barnard, W. A. Womack, S. M. Smith, T. D. Engerson, A. E. Taylor. Lung protection against paraquat is calcium dependent. <i>J Appl Physiol (1985).</i> 1992. 72:498-504
L. S. He, S. W. Chang, P. Ortiz de Montellano, T. J. Burke, N. F. Voelkel. Lung injury in Fischer but not Sprague-Dawley rats after short-term hyperoxia. <i>Am J Physiol.</i> 1990. 259:L451-8
C. Giulivi, C. C. Lavagno, F. Lucesoli, M. J. Bermudez, A. Boveris. Lung damage in paraquat poisoning and hyperbaric oxygen exposure: superoxide-mediated inhibition of phospholipase A2. <i>Free Radic Biol Med.</i> 1995. 18:203-13
W. H. Cheng, X. Zheng, F. R. Quimby, C. A. Roneker, X. G. Lei. Low levels of glutathione peroxidase 1 activity in selenium-deficient mouse liver affect c-Jun N-terminal kinase activation and p53 phosphorylation on Ser-15 in pro-oxidant-induced apoptosis. <i>Biochem J.</i> 2003. 370:927-34
J. F. Turrens, C. Giulivi, C. Pinus, E. Roldan, C. Lavagno, A. Boveris. Low level chemiluminescence from isolated rat hepatocytes, intact lung and intestine in situ. <i>Basic Life Sci.</i> 1988. 49:239-42
R. F. Burk, R. A. Lawrence, J. M. Lane. Liver necrosis and lipid peroxidation in the rat as the result of paraquat and diquat administration. Effect of selenium deficiency. <i>J Clin Invest.</i> 1980. 65:1024-31
S. Z. Cagen, J. E. Gibson. Liver damage following paraquat in selenium-deficient and diethyl maleate-pretreated mice. <i>Toxicol Appl Pharmacol.</i> 1977. 40:193-200
Z. E. Suntres, P. N. Shek. Liposomal alpha-tocopherol alleviates the progression of paraquat-induced lung damage. <i>J Drug Target.</i> 1995. 2:493-500
J. S. Bus, S. D. Aust, J. E. Gibson. Lipid peroxidation: a possible mechanism for paraquat toxicity. <i>Res Commun Chem Pathol Pharmacol.</i> 1975. 11:31-8
T. Matsumoto, H. Matsumori, Y. Hosokawa, H. Abe, T. Uekusa, N. Kuwabara, Y. Fukuda, H. Ishikawa. A light and electron microscopic study of the regenerative epithelium in the intrahepatic bile duct. Experimental study of local direct instillation of paraquat into the intrahepatic bile ducts of rats. <i>Acta Pathol Jpn.</i> 1986. 36:1163-80

Paraquat (PQ) poisoning often leads to severe oxidative liver injury. Recent studies have reported that methylene blue (M
Paraquat (PQ) is a widely used herbicide, and lung is the primary target of PQ poisoning. Metallothionein (MT) is a poten
We employed a low-molecular-weight metalloporphyrin superoxide dismutase mimetic, MnTBAP, to protect mice against
The effects of acute paraquat exposure on mitochondrial function in rat lung were studied. The paraquat dose-response
The production of superoxide anion (O ₂ ⁻) statistically increased, stimulated by paraquat (1,1'-dimethyl-4,4'-bipyridylium c
Experimental studies have shown that toxicant responsive genes, cytochrome P450s (CYPs) and glutathione S-transferase
BACKGROUND: Paraquat (PQ), an effective and widely used herbicide, has been proven to be safe when appropriately ap
Isolated, perfused, and ventilated rat lungs were challenged by paraquat (0.01 M) in the presence of 2.5 mM Ca ²⁺ , 2.5 m
The Fischer rat is known for its susceptibility to develop liver necrosis when challenged with paraquat (Smith et al., J. Pha
Paraquat poisoning and hyperbaric oxygen exposure are well established models of oxidative stress in lung. The aim of th
Low levels of hepatic selenium (Se)-dependent glutathione peroxidase 1 (GPX1) activity have been shown to protect aga
Paraquat and diquat facilitate formation of superoxide anion in biological systems, and lipid peroxidation has been postu
The present study was carried out to investigate the efficacy of liposome-associated alpha-tocopherol in treating pulmon
The toxicity of the herbicide paraquat in mice, measured by the single dose lethal to 50 per cent of the animals after 7 da
To date, no reliable report on the regeneration of the intrahepatic bile duct epithelium following damage to the duct has

Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant

Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1



1529
1530
1532
1534
1535
1536
1538
1539
1541
1542
1543
1544
1546
1547
1548
1549
1550
1551

L. Atzori, B. Cannas, T. Dettori, M. Dore, C. Montaldo, G. Ugazio, L. Congiu. Involvement of capsaicin-sensitive nerves in paraquat-induced mortality. <i>Chem Biol Interact.</i> 1998. 116:93-103
Z. E. Suntres, P. N. Shek. Intratracheally administered liposomal alpha-tocopherol protects the lung against long-term toxic effects of paraquat. <i>Biomed Environ Sci.</i> 1995. 8:289-300
M. Hu, W. Wu, J. Gong, Y. Li. [The interventional effect of Xuebijing injection on expression of mitochondrial fusion protein 2 and the ultrastructure changes in lung tissues in rats with paraquat poisoning]. <i>Zhonghua Wei Zhong Bing Ji Jiu Yi Xue.</i> 2014. 26:388-93
J. H. Cho, D. K. Yang, L. Kim, J. S. Ryu, H. L. Lee, C. M. Lim, Y. S. Koh. Inhaled nitric oxide improves the survival of the paraquat-injured rats. <i>Vascul Pharmacol.</i> 2005. 42:171-8
Y. Midorikawa, H. Ogata. [Influence of nitric oxide on superoxide anion radical generation in vivo]. <i>Masui.</i> 1996. 45:1088-95
E. Sikora, I. Bodziarczyk. Influence of diet with kale on lipid peroxides and malondialdehyde levels in blood serum of laboratory rats over intoxication with paraquat. <i>Acta Sci Pol Technol Aliment.</i> 2013. 12:91-9
J. W. Bauman, C. Madhu, J. M. McKim, Y. Liu, C. D. Klaassen. Induction of hepatic metallothionein by paraquat. <i>Toxicol Appl Pharmacol.</i> 1992. 117:233-41
H. S. Kim, Y. Manevich, S. I. Feinstein, J. H. Pak, Y. S. Ho, A. B. Fisher. Induction of 1-cys peroxiredoxin expression by oxidative stress in lung epithelial cells. <i>Am J Physiol Lung Cell Mol Physiol.</i> 2003. 285:L363-9
F. E. Harrison, J. L. Best, M. E. Meredith, C. R. Gamlin, D. B. Borza, J. M. May. Increased expression of SVCT2 in a new mouse model raises ascorbic acid in tissues and protects against paraquat-induced oxidative damage in lung. <i>PLoS One.</i> 2012. 7:e35623
J. W. Bauman, J. Liu, Y. P. Liu, C. D. Klaassen. Increase in metallothionein produced by chemicals that induce oxidative stress. <i>Toxicol Appl Pharmacol.</i> 1991. 110:347-54
W. H. Cheng, F. W. Quimby, X. G. Lei. Impacts of glutathione peroxidase-1 knockout on the protection by injected selenium against the pro-oxidant-induced liver aponecrosis and signaling in selenium-deficient mice. <i>Free Radic Biol Med.</i> 2003. 34:918-27
M. Nagao, W. D. Zhang, Y. Itakura, M. Kobayashi, Y. Yamada, K. Yagi, T. Oono, T. Takatori. Immunohistochemical localization of paraquat in skin and eye of rat. <i>Nihon Hoigaku Zasshi.</i> 1993. 47:202-6
G. Mainwaring, F. L. Lim, K. Antrobus, C. Swain, M. Clapp, I. Kimber, G. Orphanides, J. G. Moggs. Identification of early molecular pathways affected by paraquat in rat lung. <i>Toxicology.</i> 2006. 225:157-72
M. A. Rosenbaum, K. Miyazaki, L. M. Graham. Hypercholesterolemia and oxidative stress inhibit endothelial cell healing after arterial injury. <i>J Vasc Surg.</i> 2012. 55:489-96
R. Akcilar, A. Akcilar, H. Simsek, F. E. Kocak, C. Kocak, G. Yumun, Z. Bayat. Hyperbaric oxygen treatment ameliorates lung injury in paraquat intoxicated rats. <i>Int J Clin Exp Pathol.</i> 2015. 8:13034-42
J. Wilhelm, M. Frydrychova, M. Vizek. Hydrogen peroxide in the breath of rats: the effects of hypoxia and paraquat. <i>Physiol Res.</i> 1999. 48:445-9
Z. Hami, M. Amini, A. Kiani, M. Ghazi-Khansari. High Performance Liquid Chromatography Coupled with Pre-column Derivatization for Determination of Oxidized Glutathione Level in Rats Exposed to Paraquat. <i>Iran J Pharm Res.</i> 2013. 12:911-6
W. H. Cheng, B. A. Valentine, X. G. Lei. High levels of dietary vitamin E do not replace cellular glutathione peroxidase in protecting mice from acute oxidative stress. <i>J Nutr.</i> 1999. 129:1951-7

Paraquat (PQ), a broad spectrum herbicide, produces severe lung inflammation and necrosis resulting in pulmonary fibrosis.
Paraquat is a broad-spectrum herbicide known to produce lung injury via oxidative stress-mediated mechanisms. Different mechanisms have been proposed for the lung injury induced by paraquat.
OBJECTIVE: To investigate the mechanism of pulmonary fibrosis induced by paraquat (PQ), and the effect of Xuebijing injection on the paraquat-induced lung injury.
The purpose of this study was to evaluate the effects of inhaled nitric oxide (NO) on the paraquat-induced lung injury in rats.
We studied the effect of NO on superoxide anion radical (O ₂ ⁻) generation using chemiluminescence method by examining the effect of NO on the superoxide anion radical (O ₂ ⁻) generation.
BACKGROUND: Organism's lipid peroxidation is one of the most often examined and known physiological process evoked by oxidative stress.
Paraquat, a frequently used contact herbicide, produces oxidative stress by undergoing redox cycling and generating reactive oxygen species.
1-Cys peroxiredoxin (1-cysPrx), a member of the peroxiredoxin family that contains a single conserved cysteine residue, is a major antioxidant enzyme in the cytosol.
A new transgenic mouse model for global increases in the Sodium Dependent Vitamin C transporter 2 (SVCT2) has been generated.
Metallothionein (MT) is a low-molecular-weight protein with a high cysteine content that has been proposed to play a role in the detoxification of heavy metals.
Previous research has suggested that depletion of cellular glutathione peroxidase (GPX1) activity by a single injection of S-nitrosoglutathione (GSNO) can lead to increased oxidative stress.
Immunohistochemical techniques were used to observe the localization of paraquat in the skin and eyes of rats. Paraquat was found to be localized in the skin and eyes of rats.
We have used global gene expression profiling, combined with pathway analysis tools, to identify in rats the molecular events that occur during paraquat-induced lung injury.
OBJECTIVE: Endothelial cell (EC) migration is essential for arterial healing after angioplasty. Oxidized low-density lipoprotein (oxLDL) is a major factor in the development of atherosclerosis.
Paraquat (PQ) is an agrochemical agent commonly used worldwide, which can cause acute lung injury (ALI) and death. However, the mechanism of PQ-induced ALI is not fully understood.
The hypothesis that oxidative stress can be induced by hypoxia was tested by measuring the concentration of hydrogen peroxide (H ₂ O ₂) in the lungs of rats.
Glutathione (GSH) is one of the most important antioxidants that plays an essential role in detoxification of reactive oxygen species.
Our objective was to determine whether high levels of dietary vitamin E replaced the protection of the Se-dependent cell antioxidant enzymes.

Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant